

## CHAPTER 10

### OTHER FLATFISH

by

Thomas K. Wilderbuer, Daniel G. Nichol and Paul D. Spencer

#### **EXECUTIVE SUMMARY**

The following changes have been made to this assessment relative to the November 2006 SAFE:

##### Changes in the input data

- 1) The 2007 catch (total and discarded) was updated, and catch through 8 September, 2007 were included in the assessment.
- 2) 2007 Eastern Bering Sea trawl survey biomass estimates and standard errors of other flatfish species are included in the assessment. A linear regression of the AI survey estimates was used to predict the AI biomass in 2007, when an AI survey did not occur.

|   | 2007 Assessment<br>recommendations | 2006 Assessment<br>recommendations |
|---|------------------------------------|------------------------------------|
| Exploitable biomass                         | 149,497 t                          | 149,292 t                          |
| ABC   | 21,591 t                           | 21,396 t                           |
| Overfishing                                 | 28,788 t                           | 28,528 t                           |
| $F_{ABC}$ for rex sole                      | 0.13                               | 0.13                               |
| $F_{overfishing}$ for rex sole              | 0.17                               | 0.17                               |
| $F_{ABC}$ for Dover sole                    | 0.064                              | 0.064                              |
| $F_{overfishing}$ for Dover sole            | 0.085                              | 0.085                              |
| $F_{ABC}$ for the remaining species         | 0.20                               | 0.20                               |
| $F_{overfishing}$ for the remaining species | 0.15                               | 0.15                               |

## Response to SSC comments

**The SSC encourages explorations of bottom temperature relationships with catchability for the other species in the other flatfish complex. It would be useful to be able to identify indicator species that are particularly sensitive to changes in environmental conditions, and also in scaling biomass estimates appropriately for different species in the complex.**

No progress was made on these types of explorations in 2007.

**The SSC notes the catch estimates in the BSAI plan team introduction on page 20 mislabels the catch and ABC columns for 2005 and 2006, creating a corresponding error in the text. The TAC was exceeded by 31% in 2004 and 56% in 2005. The TAC was not exceeded in 2006.**

Preliminary specified TACs for “other flatfish” are usually a balance between the available Pollock and Pacific cod TACs and the 2.0 million metric t groundfish cap. In 2004 and 2005 there was not enough room under the cap to provide a larger TAC for “other flatfish” and still fulfill the desires of the fishing industry. Thus the annual catch exceeded the TAC when these species were captured, primarily in the pursuit of other flatfish species. However, in neither year did the catch exceed the ABC.

**Although biomass estimates for the dominant species are increasing, the harvest of the remaining species category listed in Table 10.2 was the lowest in the series for 2005 and 2006. The SSC requests that the “remaining species” category in Table 10.2 be speciated to track relative changes in catch, and that all “other flatfish” species in the survey data be listed individually in Table 10.5.**

The requested changes were made to Tables 10.2 and 10.5.

## Introduction

The Bering Sea/Aleutian Islands “other flatfish” group have typically included those flatfish besides rock sole, yellowfin sole, arrowtooth flounder, and Greenland turbot. Flathead sole (*Hippoglossoides elassodon*) were part of the other flatfish complex until they were removed in 1995, and Alaska plaice was removed from the complex in 2002, as sufficient biological data exists for these species to construct age-structured population models. In contrast, survey biomass estimates are the principal data source used to assess the remaining other flatfish. Although over a dozen species (Table 10.1) of flatfish are found in the BSAI area, the other flatfish biomass consists primarily of starry flounder, rex sole, longhead dab, Dover sole and butter sole.

## Catch History

The miscellaneous species of the other flatfish species category are listed in Table 10.1, and their catches from 1995-2007 are shown in Table 10.2. These species are not pursued as fishery targets but are captured in fisheries for other species. Catch from 1995-2003 were obtained from the NMFS Regional Office “blend” data, and the catch for some species are reported by species and in an aggregate flatfish group. The catch estimates for these years were produced by applying the proportional catch, by species, from fishery observer data to the estimated total catch for the aggregate other flatfish group, and adding this total to the catch that was reported by species. In the newer catch accounting system (in use since 2003), catches of other flatfish are reported only in an aggregate group, and the catch estimates for these years were produced by applying the proportional catch, by species, from fishery observer data to the estimated total catch of the aggregate group. In recent years, starry flounder (*Platichthys stellatus*) and rex sole (*Glyptocephalus zachirus*) account for most of the harvest of other flatfish, and contributed 88% of the harvest of other flatfish in 2007.

Other flatfish are grouped with Alaska plaice, rock sole, and flathead sole and other flatfish fisheries in a single prohibited species group (PSC) classification, with seasonal and total annual allowances of prohibited bycatch applied to the group. In recent years, this group of fisheries has been closed prior to attainment of the TAC due to the bycatch of halibut (Table 10.3). In 2007 the other flatfish harvest was subject to 3 closures: the first and second seasonal closures to prohibit further halibut retention, and the annual halibut cap closure as detailed in Table 10.3.

## DATA

### Absolute Abundance and Exploitation Rates

The biomass of the other flatfish complex on the eastern Bering Sea shelf has been relatively stable from 1983-1995, averaging 50,200 t, and has slightly increased from 1996 to 2005, averaging 84,500 t (Table 10.4). The 2007 biomass estimate of 133,491 t of other flatfish on the EBS shelf is the highest estimate since surveys began in 1982 and the 2006 Aleutian Islands trawl survey biomass estimate of 16,440 is the highest observed there as well. The increases are primarily due to the higher estimate of starry flounder on the Eastern Bering Sea shelf. An estimate of total BSAI biomass for the years in which an AI survey was not conducted (as in

2007) was calculated by regressing the AI survey biomass against the EBS survey biomass and adding the predicted AI biomass estimate to the observed EBS estimate. Individual species biomass estimates for the EBS and AI areas from 1997-2007 are shown in Table 10.5. Estimates of species biomass for starry flounder, rex sole, and butter sole in the Aleutian Islands were computed by fitting a linear trend to the observed survey data from 1991-2006, and using this trend to estimate biomass in years without an Aleutian Island survey. Estimate of total BSAI biomass (Table 10.6) were then used to compute species-specific exploitation rates.

Exploitation rates for starry flounder and rex sole have been low, not exceeding 0.10 from 1997 to 2007 (Table 10.6). The exploitation rates for butter sole have been higher, exceeding 0.14 in 1997, 2000, 2001, and 2003-2007, but the biomass estimates from which these were calculated for butter sole have large sampling variances, with coefficients of variation ranging from 0.44 to 0.86 in recent EBS trawl surveys dating back to 1999.

The 2003 biomass estimate of butter sole of 429 t is less than one-fourth the 2002 estimate of 2382, and results in an estimated exploitation rate of nearly 70%. However, butter sole were only captured in four hauls in the 2003 EBS trawl survey, leading to the large coefficient of variation of 0.61 for the estimated biomass. In addition, the bulk of the 2003 fishery catch came primarily from waters less than 50 m in January and February, a depth and time not covered by the trawl survey. Thus, it is likely that the population of butter sole is larger than that indicated from the survey, and the comparison of survey biomass to harvest should be interpreted accordingly. The 2007 biomass estimate of butter sole was 1104 t, between four and five times the 2003 estimate, with a high CV in both the shelf survey (0.53) and in the Aleutian Islands (0.98).

Several species of other flatfish are relatively rare on the EBS shelf, including Dover sole, Sakhalin sole, and English sole, and it is useful to identify whether the EBS represents the edge of the distribution for these species. The distribution of English sole has been identified as Baja California to Unimak Island, and the distribution of Dover sole has been identified as from Baja California to the Bering Sea (Hart 1973). Thus, the eastern Bering Sea can be considered the periphery of the range for these species. They are much more abundant in the Gulf of Alaska. For example, the abundance of Dover sole in the 1984-2001 GOA surveys has fluctuated between 63,000 t and 96,000 t, the abundance of butter sole has fluctuated between 17,000 t and 30,000 t, and the abundance of English sole has fluctuated between 3,000 t and 14,000 t (Turnock et al. 2005). Dover sole and English sole were most common in the eastern portion of the GOA, consistent with their reported distribution along the west coast of North America. In the case of Sakhalin sole, which prefer colder water and are caught at the northern extent of the survey, their perceived abundance from survey biomass estimates may be related to annual mean bottom water temperature (Fig 10.1).

## PROJECTIONS AND HARVEST ALTERNATIVES

### Reference Fishing Mortality Rates and Yields

Other flatfish are assessed under Tier 5 of Amendment 56 to the BSAI groundfish management plan, and thus have harvest recommendations which are directly calculated from estimates of

biomass and natural mortality. The natural mortality rates used in age-structured BSAI flatfish assessments can be used as guidance and are presented below:

| <u>Species</u>      | <u>Natural mortality rate used for stock assessment</u> |
|---------------------|---|
| BSAI Yellowfin sole | 0.12  |
| BSAI Rock sole      | 0.15  |
| BSAI Flathead sole  | 0.20  |
| BSAI Alaska plaice  | 0.25  |
| GOA Rex sole        | 0.17  |
| GOA Dover sole      | 0.085   |

Natural mortality values for rex and Dover sole are available from age-structured assessments in the Gulf of Alaska SAFE document (Turnock and A'mar 2005 and Stockhausen et al. 2005) and those published values are used for rex and Dover sole in this stock assessment. For the remaining flatfish species, where less information is available, an assumption of  $M = 0.2$  appears reasonable given the range of values shown above. For the case of starry flounder where estimates are available from a west coast stock assessment (Ralston 2005), the high estimates of  $M$  (male = 0.45, female = 0.3) are not used here due to the uncertainty of the estimates and the large spatial difference between the two management areas.

The estimates of  $F_{abc}$  and  $F_{ofl}$  under tier 5 are  $0.75M$  and  $M$ , respectively, and the ABC and OFL levels are the product of the fishing mortality rate and the biomass estimate. Given the  $F_{abc}$  and  $F_{ofl}$  levels of 0.15 and 0.20, and the biomass estimate of 149,497 t, the resulting ABC and OFL levels are 21,591 and 28,788 t.

|                             | <b>F<sub>ABC</sub></b> | <b>F<sub>OFL</sub></b> | <b>ABC</b>    | <b>OFL</b>    |
|-----------------------------|------------------------|------------------------|---------------|---------------|
| Rex sole                    | 0.13                   | 0.17                   | 3,959         | 5,279         |
| Dover sole                  | 0.064                  | 0.085                  | 100           | 133           |
| Others                      | 0.15                   | 0.20                   | 17,532        | 23,377        |
| <b>Total Other flatfish</b> |                        |                        | <b>21,591</b> | <b>28,788</b> |

### Summary

In summary, several quantities pertinent to the management of the other flatfish are listed below.

-

| <u>Quantity</u>         | <u>Value</u> |
|-------------------------|--------------|
| Tier                    | 5            |
| Year 2007 Total Biomass | 149,497 t    |
| OFL                     | 28,788 t     |
| Maximum allowable ABC   | 21,591 t     |
| Recommended ABC         | 21,591 t     |

## REFERENCES

- Hart, J.L. 1973. Pacific fishes of Canada. Fisheries Research Board of Canada, Bulletin 180, Ottawa. 740 pp.
- Ralston, S. 2005. Starry flounder. An assessment of starry flounder off California, Oregon and Washington. In Status of the Pacific coast groundfish fisheries through 2005. Stock assessment and fishery evaluation. Pacific Fishery Management Council, Portland Oregon.
- Spencer, P.D., T.K. Wilderbuer, and C.I. Zhang. 2002. A mixed-species yield per recruit model for eastern Bering Sea flatfish fisheries. *Can J. Fish. Aquat. Sci.* 59:291-302.
- Stockhausen, W.T., B. J. Turnock, A. T. A'mar, M. E. Wilkins and M. H. Martin. 2005. Gulf of Alaska Dover Sole. In Stock Assessment and Fishery Evaluation Document for Groundfish Resources in the Gulf of Alaska Region as Projected for 2002. North Pacific Fishery Management Council, P.O. Box 103136, Anchorage Alaska 99510.
- Turnock, B.J., T.K. Wilderbuer, and E.S. Brown. 2001. Gulf of Alaska flatfish. In Stock Assessment and Fishery Evaluation Document for Groundfish Resources in the Gulf of Alaska Region as Projected for 2006. North Pacific Fishery Management Council, P.O. Box 103136, Anchorage Alaska 99510.
- Turnock, B.J. and Z. T. A'mar. 2005. Gulf of Alaska rex sole stock assessment. In Stock Assessment and Fishery Evaluation Document for Groundfish Resources in the Gulf of Alaska Region as Projected for 2006. North Pacific Fishery Management Council, P.O. Box 103136, Anchorage Alaska 99510.

Table 10.1. Flatfish species of the Bering Sea/Aleutian Islands “other flatfish” management complex.

| <u>Common Name</u>   | <u>Scientific Name</u>                   |
|----------------------|--|
| Arctic flounder      | <i>Liopsetta glacialis</i>               |
| butter sole          | <i>Isopsetta isolepis</i>                |
| curlfin sole         | <i>Pleuronectes decurrens</i>            |
| deepsea sole         | <i>Embassichthys bathybus</i>            |
| Dover sole           | <i>Microstomus pacificus</i>             |
| English sole         | <i>Parophrys vetulus</i>                 |
| longhead dab         | <i>Limanda proboscidea</i>               |
| Pacific sanddab      | <i>Citharichthys sordidus</i>            |
| petrale sole         | <i>Eopsetta jordani</i>                  |
| rex sole             | <i>Glyptocephalus zachirus</i>           |
| roughscale sole      | <i>Clidodoerma asperrimum</i>            |
| sand sole            | <i>Psettichthys melanostictus</i>        |
| slender sole         | <i>Lyopsetta exilis</i>                  |
| starry flounder      | <i>Platichthys stellatus</i>             |
| <u>Sakhalin sole</u> | <u><i>Pleuronectes sakhalinensis</i></u> |

Table 10.2. Harvest (t) of other flatfish from 1995-2007. 2007 catch is through September 8, 2007.

| Year | Starry<br>Founder | Rex<br>Sole | Butter<br>Sole | longhead<br>dab | Dover<br>sole | English<br>sole | deep sea<br>sole | Sakhalin<br>sole | Total | ABC    | TAC    |
|------|-------------------|-------------|----------------|-----------------|---------------|-----------------|------------------|------------------|-------|--------|--------|
| 1995 | 398               | 673         | 157            | 7               | 59            | 26              | 4                | 0                | 1324  | 117000 | 19540  |
| 1996 | 1171              | 1148        | 218            | 175             | 6             | 0               | 0                | 30               | 2748  | 102000 | 35000  |
| 1997 | 1043              | 687         | 448            | 211             | 53            | 0               | 29               | 6                | 2490  | 97500  | 50750  |
| 1998 | 402               | 998         | 229            | 93              | 41            | 0               | 0                | 0                | 1765  | 164000 | 89434  |
| 1999 | 725               | 998         | 230            | 56              | 81            | 27              | 0                | 0                | 2117  | 154000 | 154000 |
| 2000 | 1151              | 1069        | 458            | 277             | 66            | 4               | 0                | 0                | 3027  | 117000 | 83813  |
| 2001 | 755               | 869         | 244            | 62              | 70            | 4               | 6                | 0                | 2028  | 122000 | 28000  |
| 2002 | 1075              | 1192        | 222            | 107             | 34            | 0               | 1                | 0                | 2631  | 18100  | 3000   |
| 2003 | 887               | 1399        | 296            | 125             | 39            | 2               | 0                | 0                | 2749  | 16000  | 3000   |
| 2004 | 2062              | 1858        | 514            | 146             | 82            | 6               | 0                | 0                | 4669  | 13500  | 3000   |
| 2005 | 2069              | 2001        | 487            | 25              | 16            | 1               | 0                | 0                | 4599  | 21400  | 3500   |
| 2006 | 1663              | 1266        | 261            | 33              | 10            | 0               | 0                | 0                | 3233  | 18100  | 3500   |
| 2007 | 4080              | 760         | 542            | 82              | 4             | 2               | 0                | 0                | 5470  | 21400  | 10000  |

Table 10.3. Restrictions on the “other flatfish” fishery from 1995 to 2006 in the Bering Sea – Aleutian Islands management area. Note that in 1994, the other flatfish category included flathead sole. Unless otherwise indicated, the closures were applied to the entire BSAI management area. Zone 1 consists of areas 508, 509, 512, and 516, whereas zone 2 consists of areas 513, 517, and 521.

| Year | Dates        | Bycatch Closure                        |
|------|--------------|--|
| 1995 | 2/21 – 3/30  | First Seasonal halibut cap             |
|      | 4/17 – 7/1   | Second seasonal halibut cap            |
|      | 8/1 – 12/31  | Annual halibut allowance               |
| 1996 | 2/26 – 4/1   | First Seasonal halibut cap             |
|      | 4/13 – 7/1   | Second seasonal halibut cap            |
|      | 7/31 – 12/31 | Annual halibut allowance               |
| 1997 | 2/20 – 4/1   | First Seasonal halibut cap             |
|      | 4/12 – 7/1   | Second seasonal halibut cap            |
|      | 7/25 – 12/31 | Annual halibut allowance               |
| 1998 | 3/5 – 3/30   | First Seasonal halibut cap             |
|      | 4/21 – 7/1   | Second seasonal halibut cap            |
|      | 8/16 – 12/31 | Annual halibut allowance               |
| 1999 | 2/26 – 3/30  | First Seasonal halibut cap             |
|      | 4/27 – 7/04  | Second seasonal halibut cap            |
|      | 8/31 – 12/31 | Annual halibut allowance               |
| 2000 | 3/4 – 3/31   | First Seasonal halibut cap             |
|      | 4/30 – 7/03  | Second seasonal halibut cap            |
|      | 8/25 – 12/31 | Annual halibut allowance               |
| 2001 | 3/20 – 3/31  | First Seasonal halibut cap             |
|      | 4/27 – 7/01  | Second seasonal halibut cap            |
|      | 8/24 – 12/31 | Annual halibut allowance               |
| 2002 | 2/22 – 12/31 | Red King crab cap (Zone 1 closed)      |
|      | 3/1 – 3/31   | First Seasonal halibut cap             |
|      | 4/20 – 6/29  | Second seasonal halibut cap            |
|      | 7/29 – 12/31 | Annual halibut allowance               |
| 2003 | 2/18 – 3/31  | First Seasonal halibut cap             |
|      | 4/1 – 6/21   | Second seasonal halibut cap            |
|      | 7/31 – 12/31 | Annual halibut allowance               |
| 2004 | 2/24 – 3/31  | First Seasonal halibut cap             |
|      | 4/10 – 12/31 | Bycatch status                         |
| 2005 | 3/1 – 3/31   | First Seasonal halibut cap             |
|      | 4/22-6/30    | Second Seasonal halibut cap            |
|      | 5/9-12/31    | Bycatch status, TAC attained           |
| 2006 | 2/21 – 3/31  | First Seasonal halibut cap             |
|      | 4/5 – 12/31  | Red King crab cap (Zone 1 closed)      |
|      | 4/12 – 5/31  | Second seasonal halibut cap            |
|      | 5/26         | TAC attained, 7,000 t reserve released |
|      | 8/7 – 12/31  | Annual halibut allowance               |
| 2007 | 2/17 – 3/31  | First Seasonal halibut cap             |
|      | 4/9 – 5/31   | Second seasonal halibut cap            |
|      | 8/6 – 12/31  | Annual halibut allowance               |

Table 10.4. Estimated biomass (t) of other flatfish from the eastern Bering Sea and Aleutian Islands trawl surveys. Species included are Dover sole, longhead dab, rex sole, Sakhalin sole, starry flounder, and butter sole. A linear regression between EBS and AI survey abundance was used to predict AI abundance in years in which an AI survey did not occur.

| Year | EBS    | Area  |         |
|------|--------|-------|---------|
|      |        | AI    | Total   |
| 1982 | 117763 |       | 129518  |
| 1983 | 66131  | 2700  | 68831   |
| 1984 | 59647  |       | 64956   |
| 1985 | 34572  |       | 37101   |
| 1986 | 39517  | 6100  | 45617   |
| 1987 | 49764  |       | 53977   |
| 1988 | 43751  |       | 47298   |
| 1989 | 49592  |       | 53786   |
| 1990 | 46649  |       | 50517   |
| 1991 | 72399  | 2144  | 74543   |
| 1992 | 53817  |       | 58480   |
| 1993 | 44399  |       | 48017   |
| 1994 | 54045  | 5464  | 59509   |
| 1995 | 37786  |       | 40671   |
| 1996 | 60225  |       | 65599   |
| 1997 | 70225  | 7580  | 77805   |
| 1998 | 73936  |       | 80830   |
| 1999 | 67713  |       | 73917   |
| 2000 | 70538  | 8149  | 78687   |
| 2001 | 78844  |       | 86282   |
| 2002 | 98052  | 8801  | 106853  |
| 2003 | 90327  |       | 99039   |
| 2004 | 127630 | 14980 | 142610  |
| 2005 | 107538 |       | 120900  |
| 2006 | 132852 | 16440 | 149292  |
| 2007 | 133491 |       | 149,497 |

Table 10.5 --Estimated biomass (t) and coefficient of variation (in parentheses) for the miscellaneous species of the “other flatfish” management complex in the Bering Sea trawl and Aleutian Islands surveys.

| <b>Eastern Bering Sea Shelf survey</b> |            |              |               |               |                 |             |              |            |
|--|------------|--------------|---------------|---------------|-----------------|-------------|--------------|------------|
| Year                                   | Dover Sole | Rex Sole     | longhead dab  | Sakhalin sole | starry flounder | butter sole | slender sole | sand sole  |
| 1982                                   | --         | 5994 (0.16)  | 103806 (0.16) | --            | 7781 (0.32)     | 182 (0.82)  | --           | --         |
| 1983                                   | --         | 7272 (0.18)  | 51386 (0.38)  | --            | 7436 (0.25)     | 37 (0.45)   | --           | 1559(0.94) |
| 1984                                   | --         | 13058 (0.28) | 35308 (0.16)  | 137 (0.43)    | 8913 (0.36)     | 2231 (0.64) | --           | --         |
| 1985                                   | 10 (1.04)  | 10751 (0.20) | 9107 (0.13)   | 102 (0.37)    | 12181 (0.24)    | 2421 (0.83) | --           | --         |
| 1986                                   | 15 (1.00)  | 12886 (0.22) | 10889 (0.14)  | 274 (0.48)    | 9112 (0.33)     | 6341 (0.58) | --           | --         |
| 1987                                   | 81 (0.91)  | 12931 (0.19) | 11897 (0.19)  | 110 (0.59)    | 22702 (0.63)    | 2043 (0.38) | --           | --         |
| 1988                                   | 38 (0.59)  | 15445 (0.15) | 16710 (0.19)  | 253 (0.63)    | 9222 (0.30)     | 2083 (0.47) | --           | 1128(1.0)  |
| 1989                                   | --         | 12939 (0.15) | 13086 (0.16)  | 58 (0.57)     | 22205 (0.35)    | 1304 (0.54) | --           | --         |
| 1990                                   | 47 (0.58)  | 11857 (0.21) | 18601 (0.15)  | 110 (0.51)    | 15048 (0.26)    | 986 (0.60)  | --           | --         |
| 1991                                   | 55 (0.70)  | 16014 (0.28) | 18680 (0.14)  | 291 (0.79)    | 34303 (0.23)    | 3056 (0.50) | --           | --         |
| 1992                                   | 137 (0.58) | 14001 (0.24) | 10827 (0.17)  | 75 (0.48)     | 27544 (0.22)    | 1233 (0.70) | --           | --         |
| 1993                                   | 37 (0.75)  | 14567 (0.32) | 11690 (0.21)  | 78 (0.34)     | 16510 (0.22)    | 1517 (0.75) | --           | --         |
| 1994                                   | 73 (0.72)  | 15943 (0.38) | 18533 (0.26)  | 183 (0.41)    | 18218 (0.22)    | 1095 (0.97) | --           | --         |
| 1995                                   | --         | 10420 (0.28) | 8402 (0.15)   | 109 (0.32)    | 17652 (0.29)    | 1203 (0.54) | --           | --         |
| 1996                                   | --         | 10532 (0.40) | 8567 (0.20)   | 34 (0.34)     | 40409 (0.45)    | 683 (0.53)  | --           | --         |
| 1997                                   | --         | 8233 (0.27)  | 18003 (0.21)  | 87 (0.49)     | 41018 (0.21)    | 2884 (0.43) | --           | --         |
| 1998                                   | 41 (0.44)  | 7588 (0.22)  | 14737 (0.19)  | 34 (0.49)     | 49605 (0.30)    | 1942 (0.38) | --           | --         |
| 1999                                   | 16 (0.65)  | 8020 (0.28)  | 12087 (0.21)  | 63 (0.29)     | 43375 (0.25)    | 4152 (0.62) | --           | --         |
| 2000                                   | 11 (1.02)  | 9348 (0.19)  | 13511 (0.30)  | 145 (0.88)    | 45810 (0.19)    | 1713 (0.56) | --           | --         |
| 2001                                   | 16 (0.84)  | 21660 (0.23) | 12764 (0.26)  | 31 (0.43)     | 43026 (0.25)    | 796 (0.50)  | --           | --         |
| 2002                                   | 7 (0.80)   | 26053 (0.20) | 9740 (0.22)   | 7 (0.69)      | 59877 (0.23)    | 2254 (0.64) | --           | --         |
| 2003                                   | 350 (0.66) | 28023 (0.15) | 8827(0.22)    | 55 (0.40)     | 52893 (0.17)    | 179 (0.61)  | 3            | --         |
| 2004                                   | 31(0.51)   | 28762 (0.19) | 11290 (0.23)  | 8 (0.64)      | 86698 (0.38)    | 841 (0.86)  | --           | --         |
| 2005                                   | 157(0.19)  | 23171(0.19)  | 11556 (0.21)  | 23(0.90)      | 71673(0.26)     | 958(0.81)   | --           | --         |
| 2006                                   | 90(0.53)   | 21515(0.28)  | 13204(0.25)   | 52(0.41)      | 96900(0.37)     | 1091(0.53)  | --           | --         |
| 2007                                   | 73(0.53)   | 17025(0.25)  | 16733(0.24)   | 19(0.40)      | 98623(0.17)     | 1018(0.44)  | --           | --         |

**Aleutian Islands Surveys**

| Year | Species     |              |              |               |                 |             |              |
|------|-------------|--------------|--------------|---------------|-----------------|-------------|--------------|
|      | Dover Sole  | Rex Sole     | longhead dab | Sakhalin sole | starry flounder | butter sole | English sole |
| 1991 | 174 (0.45)  | 1694 (0.18)  | --           | --            | 142 (0.85)      | 86 (0.73)   | 47 (0.80)    |
| 1994 | 438 (0.41)  | 4306 (0.15)  | --           | --            | 134 (0.69)      | 505 (0.98)  | 83 (0.81)    |
| 1997 | 386 (0.34)  | 6378 (0.16)  | --           | --            | 459 (0.90)      | 346 (0.98)  | 12 (0.72)    |
| 2000 | 630 (0.38)  | 6526 (0.18)  | --           | --            | 590 (0.71)      | 310 (0.99)  | 95 (0.97)    |
| 2002 | 575 (0.28)  | 7381 (0.15)  | --           | --            | 671 (0.72)      | 127 (0.83)  | 47 (0.94)    |
| 2004 | 870 (0.28)  | 13717 (0.18) | --           | --            | 123 (0.72)      | 235 (0.93)  | 35(1.00)     |
| 2006 | 2155 (0.57) | 14230 (0.19) | --           | --            | 17 (0.97)       | 13(0.98)    | 25(0.84)     |

Table 10.6. Estimated exploitation rates of rex sole, starry flounder and butter sole from 1997 to 2007.

| Year | Rex sole    |             |           | Starry Flounder |             |           | Butter sole |             |           |
|------|-------------|-------------|-----------|-----------------|-------------|-----------|-------------|-------------|-----------|
|      | Biomass (t) | Harvest (t) | Exp. Rate | Biomass (t)     | Harvest (t) | Exp. Rate | Biomass (t) | Harvest (t) | Exp. Rate |
| 1997 | 14611       | 401         | 0.03      | 41477           | 814         | 0.02      | 3230        | 336         | 0.10      |
| 1998 | 14250       | 569         | 0.04      | 49950           | 242         | 0.00      | 2210        | 157         | 0.07      |
| 1999 | 15415       | 516         | 0.03      | 43750           | 597         | 0.01      | 4416        | 167         | 0.04      |
| 2000 | 15874       | 569         | 0.04      | 46400           | 770         | 0.02      | 2023        | 266         | 0.13      |
| 2001 | 30524       | 507         | 0.02      | 43829           | 479         | 0.01      | 1059        | 147         | 0.14      |
| 2002 | 33411       | 1227        | 0.04      | 60633           | 1023        | 0.02      | 2382        | 187         | 0.08      |
| 2003 | 38349       | 1399        | 0.04      | 53353           | 887         | 0.02      | 429         | 296         | 0.69      |
| 2004 | 42479       | 1858        | 0.04      | 86821           | 2062        | 0.02      | 1076        | 514         | 0.48      |
| 2005 | 34963       | 1830        | 0.05      | 72176           | 1892        | 0.03      | 1201        | 445         | 0.37      |
| 2006 | 35745       | 1266        | 0.04      | 96917           | 1663        | 0.02      | 1104        | 261         | 0.24      |
| 2007 | 17,025      |             |           | 98623           |             |           | 1018        |             |           |




Figure 10.1—Relationship between annual survey bottom water temperature anomalies (yellow bars) and Sakhalin sole biomass estimates (red line).